

Figure D1. Northeast Region Atlantic Cod – Georges Bank fishing mortality has not been controlled and biomass has not increased. Due to the periodic recalculation of F and B by stock assessment scientists, the initial estimates of F and B used in the overfished declaration are included to illustrate the uncertainty of stock assessment estimates.




Figure D2. Northeast Region White Hake – Gulf of Maine / Georges Bank fishing mortality has not been controlled and biomass has not increased. B_{msy} proxy is in kg/tow. Due to the periodic recalculation of F and B by stock assessment scientists, the initial estimates of F and B used in the overfished declaration are included to illustrate the uncertainty of stock assessment estimates.




Figure D3. Northeast Region Yellowtail Flounder – Cape Cod / Gulf of Maine fishing mortality has not been controlled and biomass has not increased. Due to the periodic recalculation of F and B by stock assessment scientists, the initial estimates of F and B used in the overfished declaration are included to illustrate the uncertainty of stock assessment estimates.




Figure D4. Northeast Region Yellowtail Flounder – Southern New England / Mid-Atlantic fishing mortality has not been controlled and biomass has not increased. Due to the periodic recalculation of F and B by stock assessment scientists, the initial estimates of F and B used in the overfished declaration are included to illustrate the uncertainty of stock assessment estimates.




Figure D5. Blue Marlin - Atlantic fishing mortality has not been controlled and biomass has not increased. *FMP has not been internationally implemented.




Figure D5. Dusky Shark - Atlantic




Figure D7. Southeast Region Greater Amberjack – Gulf of Mexico fishing mortality is not controlled and biomass has not increased.